

Journal of Health Monitoring · 2017 2(3)
DOI 10.17886/RKI-GBE-2017-069
Robert Koch Institute, Berlin

Authors:

Julia Thom, Ronny Kuhnert,
Sabine Born, Ulfert Hapke

12-month prevalence of self-reported medical diagnoses of depression in Germany

Abstract

Depression is a frequent mental disorder and has a growing importance in health care provision. In GEDA 2014/2015-EHIS, 9.7% of women and 6.3% of men self-reported having received a medical diagnosis of depression during the past 12 months. For both genders, the rate of self-reported diagnoses of depression is highest in the 45- to 64-year age group. Education thereby plays a significant role. Prevalence for women from low education is about double that of women from high education backgrounds (12.2% compared with 6.5%). The education gradient for men is smaller (7.5% compared with 5.1%). Prevalence rates also differ sharply between federal states (for women, between 5.4% and 13.4%; for men, between 3.3% and 9.4%). These results are discussed in the light of data currently available.

◆ DEPRESSION · MEDICAL DIAGNOSIS · ADULTS · HEALTH MONITORING · GERMANY

Introduction

Depression is a mental disorder that is characterised by despondency, lack of motivation, severe weariness and the loss of interest in activities that used to produce pleasure [1]. Further symptoms include difficulties concentrating, a lack of self-confidence and suicidal tendencies in more severe cases. For those affected, depression implies severe impacts on quality of life and the ability to lead a productive life [2]. Among all chronic diseases, depression accounts for the greatest number of disability-adjusted years of life [3] and is considered to be a factor in at least half of all accomplished suicides [4]. Social insurance policies document the increasing care relevance of depression and its role in cases where people become incapable of working, require rehabilitation services and/or retire [5-8]. However, based on the epide-

miological data available, the rate of depression in the population is a controversial issue [9-11]. To measure the prevalence of depression, beside further indicators, health monitoring at the Robert Koch Institute also collects data on self-reported medical diagnoses of depression.

Indicator

To survey self-reported medical diagnoses of depression, the GEDA 2014/2015-EHIS survey used self-administered paper-based and online questionnaires. Respondents were asked, 'During the past 12 months, have you had one of the following diseases or disorders?', followed by a list of diseases which also included depression. In the face of previous surveys and to increase the interpretive and comparative value of this data, the discussion in the

GEDA 2014/2015-EHIS

Data holder: Robert Koch Institute

Aims: To provide reliable information about the population's health status, health-related behaviour and health care in Germany, with the possibility of a European comparison

Method: Questionnaires completed on paper or online

Population: People aged 18 years and above with permanent residency in Germany

Sampling: Registry office sample; randomly selected individuals from 301 communities in Germany were invited to participate

Participants: 24,016 people (13,144 women; 10,872 men)

Response rate: 26.9%

Study period: November 2014 - July 2015

Data protection: This study was undertaken in strict accordance with the data protection regulations set out in the German Federal Data Protection Act and was approved by the German Federal Commissioner for Data Protection and Freedom of Information. Participation in the study was voluntary. The participants were fully informed about the study's aims and content, and about data protection. All participants provided written informed consent.

More information in German is available at www.geda-studie.de

following sections only considers respondents who said they had suffered from depression during the past 12 months and also reported having been 'diagnosed at least once by a doctor' with depression. This led to the exclusion of 26.4% (n=657) of respondents who reported depression during the past 12 months but failed to provide a lifetime medical diagnosis.

Whilst such an approach allows for efficient estimates on the prevalence of depression and is also widely used in international health surveys [12, 13], the approach is nonetheless tied to numerous prerequisites and therefore also has its limitations. Respondents need to have 1) consulted a physician; 2) received the diagnosis of depression; 3) this diagnosis needs to meet the diagnostic criteria; and 4) be reported by a physician. When taking part in the survey, the respondent moreover needs to 5) remember having received the diagnosis and 6) be willing to report the diagnosis. Furthermore, this is based on the assumption that psychological psychotherapists who offer specialist medical care and also provide diagnoses of depression are categorised as a sub-group within the larger group of physicians.

The analyses are based on data from 23,179 participants aged 18 years and older (12,777 women and 10,402 men) with valid data on self-reported medical diagnoses of depression. The calculations were carried out using a weighting factor that corrects for deviations within the sample from the German population (as of 31 December 2014) with regard to gender, age, district type and education. The district type reflects the degree of urbanisation and accounts for the regional distribution in Germany. The International Standard Classification of Edu-

cation (ISCED) was used to classify the responses provided on educational level [14]. Differences between these groups are interpreted as statistically significant if the respective confidence intervals do not overlap.

A detailed description of the methodology used in the GEDA 2014/2015-EHIS study can be found in Lange et al. 2017 [15] as well as in the article [German Health Update: New data for Germany and Europe](#), which was published in Issue 1/2017 of the Journal of Health Monitoring.

Results and discussion

This section presents the results of the analyses, discusses them in the context of further findings from health monitoring and contrasts them with an analysis of the data received from health insurance funds.

The 12-month prevalence of self-reported medical diagnoses of depression in the overall population was 8.1% (Table 1). Women (9.7%) report the diagnosis of depression significantly more often than men (6.3%). Prevalence in both genders is highest in the 45- to 64-year group. These findings confirm the known gender imbalance for mental disorders. The higher prevalence of depression among women compared with men is a classic and apparently stable epidemiological finding, a fact which is confirmed by studies that used numerous different forms of measurement, were implemented in various countries and over long periods of time [16]. Differences between the genders also exist regarding their willingness to seek help because faced with a depressive disorder women are more likely to seek therapy than men [17]. The debate on differences between the genders

Table 1
12-month prevalence of self-reported medical diagnoses of depression diagnosed by a physician according to gender, age and educational level (n=12,777 women; n=10,402 men)
Source: GEDA 2014/2015-EHIS

9.7% of women and 6.3% of men reported a medical diagnosis of depression during the past 12 months.

Women	%	(95% CI)
Women total	9.7	(9.0-10.3)
18-29 Years	8.1	(6.7-9.7)
Low education	12.3	(8.8-16.9)
Medium education	7.5	(6.0-9.4)
High education	3.6	(2.2-5.8)
30-44 Years	9.3	(8.0-10.8)
Low education	13.4	(9.3-18.9)
Medium education	10.2	(8.5-12.1)
High education	4.8	(3.6-6.4)
45-64 Years	11.8	(10.8-12.9)
Low education	15.1	(12.1-18.7)
Medium education	11.7	(10.4-13.1)
High education	9.3	(7.7-11.2)
≥ 65 Years	8.0	(6.7-9.5)
Low education	10.1	(7.9-12.8)
Medium education	6.9	(5.3-8.9)
High education	5.3	(3.4-8.1)
Total (women and men)	8.1	(7.6-8.5)

CI=Confidence interval

Men	%	(95% CI)
Men total	6.3	(5.8-6.9)
18-29 Years	4.3	(3.2-5.9)
Low education	7.0	(4.2-11.2)
Medium education	3.4	(2.3-4.9)
High education	3.8	(1.5-8.9)
30-44 Years	5.7	(4.5-7.2)
Low education	8.1	(4.5-14.1)
Medium education	6.6	(5.2-8.4)
High education	3.2	(2.1-5.0)
45-64 Years	8.5	(7.5-9.6)
Low education	9.1	(6.5-12.6)
Medium education	9.3	(7.8-11.0)
High education	7.0	(5.7-8.5)
≥ 65 Years	5.0	(4.0-6.1)
Low education	5.6	(3.7-8.3)
Medium education	5.2	(3.8-7.1)
High education	4.2	(2.9-5.9)
Total (women and men)	8.1	(7.6-8.5)

explains these facts by pointing to both biological mechanisms and the effects of gender roles as well as factors of social stress. On the other hand, these differences are also interpreted as a distortion which results from a selection of diagnostic criteria that more typically reflects female symptoms of depression and therefore under-rates depression among men [18, 19].

Increasing levels of education almost halve the prevalence of self-reported medical diagnoses of depression in the overall population (low education of 10.5% vs. high education background of 5.6%, data not shown). The education gradient in the group of women up to the age of 64 with a diagnosed depression is stronger and


statistically more relevant than for men. Besides age and gender, the year of data collection also impacts the correlation between education and the prevalence of self-reported medical diagnoses of depression [20-22]. When income and professional status are considered as factors next to education, this leads to equally inconsistent patterns [23-25].

Prevalence rates of self-reported medical diagnoses of depression vary considerably between federal states. Prevalence is highest in the city states (13.4% of women in Bremen and 9.4% of men in Berlin) (Figure 1). Prevalence in the federal states that report the lowest rates of self-reported medical diagnoses of depression is less

Figure 1
12-month prevalence of self-reported medical diagnoses of depression according to gender and federal state
(n=12,777 women; n=10,402 men)
Source: GEDA 2014/2015-EHIS



Significant differences in rates of self-reported medical diagnoses of depression exist between the federal states.



than half of this and affects 5.4% of women in Thuringia and 3.3% of men in Saxony-Anhalt. Excluding Bavaria, where prevalence is low, the map reveals an east to west gradient for men. Surveys from previous years [25] and data from health insurance funds [6, 11] evidence comparable differences between federal states. When com-

paring urban and rural areas, both of these sources of data highlight that prevalence is highest in the major cities and lowest in provincial towns [11, 23, 24]. In the accounts data from statutory health insurance funds, the frequency of depression diagnoses at the level of individual districts can vary by the factor 3 (between 5.3%

Prevalence of self-reported medical diagnoses of depression is highest at age 45 to 64.

and 18.2%) [26] and non-associated towns (between 7.2% and 21.4%) [11] even if regional differences are adjusted for age, gender and physical morbidity. Complex differences between regions must be taken into account to explain the unequal spatial distribution, such as varying concentrations of risk and protective factors, local factors that influence how willing the population is to seek help, local availability of treatment options as well as the frequency with which depression being treated is recognised and documented [26, 27]. As evidenced by a comparison with surveys from the past few years, the number of self-reported medical diagnoses of depression is no longer rising. Whereas in GEDA 2009, 8% of women and 4.5% of men reported depression [20], GEDA 2014/2015-EHIS results are comparable to the findings presented in GEDA 2012 (women 9.8%; men 6.1%) [22]. However, it has to be considered that the form of data collection has changed between these older surveys (a telephone interview) and GEDA 2014/15-EHIS (a self-administered paper-based or online questionnaire), which might have influenced responses.

This trend is also reflected in the diagnoses of depression according to the accounting data of health insurance funds. This data reveals a continuous increase in the reporting of medical classifications by physicians related to depressive disorders as a cause of incapacity to work over the past few years [6-8]. An evaluation of Company Health Insurance Fund (BKK) data reveals that depression-related absences from work more than doubled between 2003 and 2013 [8]. Using health insurance fund data to calculate values for the 12-month prevalence of depression would lead to rates between 10% and 13%

depending on individual funds [11, 26, 28, 29]. Differences between the self-reported medical diagnoses of depression published in GEDA 2014/2015-EHIS and depression diagnoses as recorded by insurance funds exist for example concerning age distribution [6, 20-23, 26, 28]. These facts indicate conceptual differences between the data collected in surveys and accounting data [30]. Accounting data for example depends highly on the capacity of doctors to provide correct medical classifications and the validity of this data on depressive disorders is questionable [29, 31]. On the other hand, the significance of the survey data presented here depends on the degree with which the survey represents the population (response bias) as well as the above-mentioned limitations regarding the indicator itself (willingness to seek help, recall and reporting bias).

Whether a self-reported medical diagnosis of depression actually indicates depression according to clinical diagnostic criteria was a question that was analysed using data from the German Health Interview and Examination Survey for Adults (DEGS1) and its additional mental health module (DEGS1-MH) [32]. Standardised clinical interviews according to current classification criteria detect depression in only 37.2% of respondents who self-reported a medical diagnosis of depression during the past 12 months. 36.2% fulfil the criteria for a different mental disorder, whereas in 26.6% of cases the diagnosis reveals no mental disorder. On the other hand, only 33.0% of those who are diagnosed with depression in a clinical interview report a medical diagnosis of depression. Estimates on the prevalence of depression therefore both under- and overestimate the number of

The 12-month prevalence of self-reported medically diagnosed depression decreases with increasing education.

cases depending on whether diagnostic criteria or self-reported medical diagnoses are used as a basis. Surveys that collect epidemiological data and link it to the data from health insurance funds could provide more differentiated results (data linkage).

If, instead of looking at the diagnoses physicians provide, we use a questionnaire to survey the presence of individual symptoms of depression during the past two weeks (PHQ-8 [33, 34]), 10.1% of the population show [depressive symptoms](#). Women are affected more often than men, there are clear regional differences and contradictory findings related to age. Whereas women aged 18 to 29 show the highest rates of depressive symptoms, it is women aged 45 to 64 who most often report a medical diagnosis of depression. For men, the number of those with depressive symptoms nearly halves at age 65 and above, but remains constant up to that age. This also highlights the fact that pressures can only be reflected in documented diagnoses once patients turn to a doctor. Compared to men, women seek medical consultation more often, a fact which is also true for younger people compared to older people [35]. Physicians who diagnose depressive symptoms in a patient do not necessarily medically classify these as depression. This is the case when for example the number and severity of depressive symptoms is low and do not therefore fulfil the general diagnosis of the disorder or when patients present further symptoms that are then collectively classified as a different mental disorder.

The data on self-reported medical diagnoses of depression published in the health survey enable, as long as we consider the limitations described, a descrip-

tion of the people who receive the diagnosis of depression in the healthcare system. The socio-demographic, socio-economic and regional imbalances in rates of diagnosis of depression reflect as many differences in morbidity as levels of care provision between different groups in the population. An analysis of data based on standardised diagnosed depression within a clinical interview compared to self-reported medical diagnoses highlights that different indicators in epidemiology and healthcare services diagnose reveal different groups of people with depression. Clarifying these discrepancies could contribute to a provision of services according to need, for example through an increased use of screening instruments in medical practice.

References

1. Dilling H, Mombour W, Schmidt MH et al. (eds) (2016) Internationale Klassifikation psychischer Störungen. ICD-10 Kapitel V (F). Diagnostische Kriterien für Forschung und Praxis. In: WHO (Hrsg). Göttingen, Hogrefe
2. Mack S, Jacobi F, Beesdo-Baum K et al. (2015) Functional disability and quality of life decrements in mental disorders: Results from the Mental Health Module of the German Health Interview and Examination Survey for Adults (DEGS1-MH). *European Psychiatry* 30:793-800
3. World Health Organization (eds) (2016) Global Health Estimates 2015: Disease burden by Cause, Age, Sex, by Country and by Region, 2000-2015. WHO, Genf
4. Hawton K, Casanas ICC, Haw C et al. (2013) Risk factors for suicide in individuals with depression: a systematic review. *J Affect Disord* 147(1-3):17-28
5. Deutsche Rentenversicherung Bund (eds) (2014) Positionspapier der Deutschen Rentenversicherung zur Bedeutung psychischer Erkrankungen in der Rehabilitation und bei Erwerbsminderung, Berlin
6. Klaber J, Günster C, Gerste B et al. (eds) (2014) Versorgungsreport 2013/2014. Schwerpunkt: Depression. Schattauer, Stuttgart
7. Techniker Krankenkasse (eds) (2015) Depressionsatlas. Auswertungen zu Arbeitsunfähigkeit und Arzneverordnungen. AQUA - Institut für angewandte Qualitätsförderung und Forschung im Gesundheitswesen GmbH, Göttingen
8. Kliner K, Rennert D, Richter M (2015) Gesundheit in Regionen – Blickpunkt Psyche. BKK Dachverband, BKK Gesundheitsatlas 2015. MWV Medizinisch Wissenschaftliche Verlagsgesellschaft, Berlin
9. Richter D, Berger K (2013) Nehmen psychische Störungen zu? *Psychiatrische Praxis* 40:176-182
10. Jacobi F, Bretschneider J, Müllender S (2015) Veränderungen und Variationen der Häufigkeit psychischer Störungen in Deutschland. Krankenkassenstatistiken und epidemiologische Befunde. In: Kliner K, Rennert D, Richter M (eds) Gesundheit in Regionen – Blickpunkt Psyche BKK Gesundheitsatlas 2015. Medizinisch wissenschaftliche Verlagsgesellschaft, Berlin, P. 63-71
11. Melchior H, Schulz H, Härter M (2014) Faktencheck Gesundheit – Regionale Unterschiede in der Diagnostik und Behandlung von Depressionen. Bertelsmann Stiftung, Gütersloh
12. Strine TW, Mokdad AH, Balluz LS et al. (2008) Depression and anxiety in the United States: findings from the 2006 Behavioral Risk Factor Surveillance System. *Psychiatr Serv* 59(12):1383-1390
13. Substance Abuse and Mental Health Services Administration (2011) Results from the 2010 National Survey on Drug Use and Health: Summary of National Findings. Substance Abuse and Mental Health Services Administration, Rockville
14. Eurostat (2016) International standard classification of education (ISCED). http://ec.europa.eu/eurostat/statistics-explained/index.php/Glossary:International_standard_classification_of_education_%28ISCED%29 (As at 13.01.2017)
15. Lange C, Finger JD, Allen J et al. (in press) Implementation of the European Health Interview Survey (EHIS) in Germany. Further development of the German Health Update (GEDA). *Archives of Public Health*
16. Parker G, Brotchie H (2010) Gender differences in depression. *International Review of Psychiatry* 22(5):429-436
17. Moller-Leimkuhler AM (2002) Barriers to help-seeking by men: a review of sociocultural and clinical literature with particular reference to depression. *J Affect Disord* 71(1-3):1-9
18. Karger A (2014) Geschlechtsspezifische Aspekte bei depressiven Erkrankungen. *Bundesgesundheitsbl - Gesundheitsforsch - Gesundheitsschutz* 57(9):1092-1098
19. Karger A (2012) Sind Männer anders depressiv? *Ärztliche Psychotherapie und Psychosomatische Medizin* 7:224-229
20. Robert Koch-Institut (eds) (2011) Daten und Fakten: Ergebnisse der Studie "Gesundheit in Deutschland aktuell 2009". Beiträge zur Gesundheitsberichterstattung des Bundes. Berlin http://edoc.rki.de/documents/rki_fv/reQXTR7OSGFRg/PDF/20dqmVRMM57G6.pdf (As at 13.07.2017)
21. Robert Koch-Institut (eds) (2012) Daten und Fakten: Ergebnisse der Studie "Gesundheit in Deutschland aktuell 2010". Beiträge zur Gesundheitsberichterstattung des Bundes. Berlin http://edoc.rki.de/documents/rki_fv/remDCCtjOJxl/PDF/21TgK-GZEOWNCY.pdf (As at 13.07.2017)
22. Robert Koch-Institut (eds) (2014) Daten und Fakten: Ergebnisse der Studie "Gesundheit in Deutschland aktuell 2012". Beiträge zur Gesundheitsberichterstattung des Bundes. Berlin http://edoc.rki.de/documents/rki_fv/recJuHnzacx8A/PDF/28Gs-WuNtFjVqY.pdf (As at 13.07.2017)

23. Busch MA, Maske UE, Ryl L et al. (2013) Prevalence of depressive symptoms and diagnosed depression among adults in Germany – Results of the German Health Interview and Examination Survey for Adults (DEGS1). Bundesgesundheitsbl - Gesundheitsforsch - Gesundheitsschutz 56(5-6):733-739
<http://edoc.rki.de/oa/articles/reLPtVNhg9T2/PDF/22QgLkX1lQ-gU.pdf> (As at 13.07.2017)
24. Maske UE, Buttery AK, Beesdo-Baum K et al. (2016) Prevalence and correlates of DSM-IV-TR major depressive disorder, self-reported diagnosed depression and current depressive symptoms among adults in Germany. Journal of Affective Disorders 190:167-177
25. Müters S, Hoebel J, Lange C (2013) Diagnosis Depression: Differences in Women and Men. Robert Koch-Institut, Berlin
<http://edoc.rki.de/series/gbe-kompakt/4-2/PDF/2.pdf> (As at 13.07.2017)
26. Erhart M, Stillfried D (2012) Analyse regionaler Unterschiede in der Prävalenz und Versorgung depressiver Störungen auf Basis vertragsärztlicher Abrechnungsdaten – Teil 1 Prävalenz. Versorgungsatlas. Zentralinstitut für die kassenärztliche Versorgung in Deutschland, Berlin
27. Melchior H, Schulz H, Härter M (2014) Stellenwert regionaler Variationen in der Prävalenz und Behandlung depressiver Erkrankungen und Implikationen für die Versorgungsforschung. Bundesgesundheitsbl - Gesundheitsforsch - Gesundheitsschutz 57:224-233
28. Grobe T, Kleine-Budde K, Bramesfeld A et al. (in preparation) Prävalenzen von Depressionen bei Erwachsenen – eine vergleichende Analyse bundesweiter Survey- und Routinedaten
29. Institut für Gesundheits- und Sozialforschung (2012) Bewertung der Kodierqualität von vertragsärztlichen Diagnosen – Eine Studie im Auftrag des GKV-Spitzenverbands in Kooperation mit der BARMER GEK. IGES Institut für Gesundheits- und Sozialforschung GmbH, Berlin
30. Geyer S, Jaunzeme J (2014) Möglichkeiten und Grenzen von Befragungsdaten und Daten gesetzlicher Krankenversicherungen. In: Swart E, Ihle P, Gothe H et al. (eds) Routinedaten im Gesundheitswesen Handbuch Sekundärdatenanalyse: Grundlagen, Methoden und Perspektiven. Verlag Hans Huber, Bern, P. 223-233
31. Gerste B, Roick C (2014) Prävalenz und Inzidenz sowie Versorgung depressiver Erkrankungen in Deutschland - Eine Analyse auf Basis der in Routinedaten dokumentierten Depressionsdiagnosen. In: Klauber J, Günster C, Gerste B et al. (eds) Versorgungs-Report 2013/2014. Schattauer, Stuttgart, P. 21-54
32. Maske UE, Hapke U, Riedel-Heller SG et al. (2017) Respondents' report of a clinician-diagnosed depression in health surveys: comparison with DSM-IV mental disorders in the general adult population in Germany. BMC Psychiatry 17(1):39
33. Kroenke K, Strine TW, Spitzer RL et al. (2009) The PHQ-8 as a measure of current depression in the general population. J Affect Disord 114(1-3):163-173
34. Löwe B, Spitzer RL, Zipfel S et al. (2002) Gesundheitsfragebogen für Patienten (PHQ-D). Kompletteversion und Kurzform. Testmappe mit Manual, Fragebögen, Schablonen. 2. Auflage. Pfizer, Karlsruhe
35. Rattay P, Butschalowsky H, Rommel A et al. (2013) Utilisation of outpatient and inpatient health services in Germany – Results of the German Health Interview and Examination Survey for Adults (DEGS1). Bundesgesundheitsbl - Gesundheitsforsch - Gesundheitsschutz 56:832-844
<http://edoc.rki.de/oa/articles/reWsjeZR5HISQ/PDF/21nRhaJo-8ha7o.pdf> (As at 13.07.2017)

Imprint

Journal of Health Monitoring

Author details

Robert Koch Institute
Department of Epidemiology and Health Monitoring, Berlin

Corresponding author
Julia Thom
Robert Koch Institute
Department of Epidemiology and Health Monitoring
General-Pape-Str. 62–66
D-12101 Berlin, Germany
E-mail: ThomJ@rki.de

Conflicts of interest

The authors declared no conflicts of interest.

Funding

The GEDA study was funded by the Robert Koch Institute and the German Federal Ministry of Health.

Note

External contributions do not necessarily reflect the opinions of the Robert Koch Institute.

Publisher

Robert Koch Institute
Nordufer 20
D-13353 Berlin, Germany

Editors

Susanne Bartig, Johanna Gutsche, Dr Franziska Prütz,
Martina Rabenberg, Alexander Rommel, Dr Anke-Christine Saß,
Stefanie Seeling, Martin Thißen, Dr Thomas Ziese
Robert Koch Institute
Department of Epidemiology and Health Monitoring
General-Pape-Str. 62–66
D-12101 Berlin
Phone: +49 (0)30-18 754-3400
E-mail: healthmonitoring@rki.de
www.rki.de/journalhealthmonitoring-en

Typesetting

Gisela Dugnus, Alexander Krönke, Kerstin Möllerke

Translation

Simon Phillips/Tim Jack

Please cite this publication as

Thom J, Kuhnert R, Born S et al. (2017) 12-month prevalence of self-reported medical diagnoses of depression in Germany. Journal of Health Monitoring 2(3):68–76
DOI 10.17886/RKI-GBE-2017-069

ISSN 2511-2708



This work is licensed under a
Creative Commons Attribution 4.0
International License.



The Robert Koch Institute is a Federal Institute within
the portfolio of the German Federal Ministry of Health